



Client: Black Canyon Veterinary Clinic  
Organization: blackcanyonvets@gmail.com  
Email: 9707872282 / 9704174946  
Phone / fax: Tammie Rendon  
Owner: Rachel Blankmeyer DVM  
Practitioner: Ranch House Locked and Loaded by Rocky Mtn  
Patient: Dog : ~~Bernedoodle~~ Poodle  
Species/Breed: 10 Months  
Age: male  
Gender: 3/5/2018 4:42:21PM  
Submitted Date: 3/5/2018 6:12:39PM  
Completed Date:

Comments: Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of the hips the hips would be rated as excellent in conformation. Radiographs obtained at two years of age will need to be obtained to completely exclude elbow and hip dysplasia and to establish a final rating for the coxofemoral joints.

### Consulting Radiologist:

Kent Wallace, DVM, MS, DACVR  
kwallace@twcny.rr.com,



Client: Black Canyon Veterinary Clinic  
Organization: blackcanyonvets@gmail.com  
Email: 9707872282 / 9704174946  
Phone / fax: Tammie Rendon  
Owner: Rachel Blankmeyer DVM  
Practitioner: Ranch House Locked and Loaded by Rocky Mtn  
Patient: Dog : ~~Bernedoodle~~ Poodle  
Species/Breed: 10 Months  
Age: male  
Gender: 3/5/2018 4:42:21PM  
Submitted Date: 3/5/2018 6:12:39PM  
Completed Date:

Case status: Completed

#### Case notes:

hips and elbows radiographs for evaluation per OFA standards

Registered name: Ranch House Locked and Loaded by Rocky Mtn  
Short name: Colt

#### The report:

412409 KW/lb

Three digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 03/05/18.

#### Radiographic Findings:

Both femoral heads are located in their respective acetabula and are congruous with the acetabular rims.

Greater than 50% of the right femoral head and approximately 50% of the left femoral head are located medial to the dorsal acetabular rims.

No osteophyte formation is present on the femoral necks or acetabula.

Alignment of the patellae with the femurs is normal.

Muscle volume in both hind legs is within normal limits.

The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation.

No sclerosis is associated with the ulnar notches.

Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

Diagnosis: Normal pelvis and elbows.